


협의되지 않은 자료의 외부제공을 금합니다.

RE:DERM PLASMA

- *NON THERMAL COLD AIR JET PLASMA*
- *BLUE LIGHT*
- *EASY OF USE*
- *FAST EFFECT*
- *NO SIDE EFFECT*
- *NO DOWN TIME*



아토피


여드름

지루성
두피염

무좀

화상

상처



[Clin Ter. 2014;165\(2\):91-3. doi: 10.7471/CT.2014.1682.](#)

High IgE levels in patients affected by psoriasis: review of the literature and personal observations.

Paparo SB¹, Guaragna MA¹, Albanesi M².

⊕ Author information


Abstract

OBJECTIVE: The concomitance of psoriasis and high IgE levels in the same patient, possibly associated to allergies, should not come as a surprise anymore, because both diseases are characterised by an immunological disorder, involving cytokines and other inflammatory mediators, with massive activation of the cell-mediated immunity.


MATERIALS AND METHODS: During a period of 18 months, using a radio-immunologic method, we assessed the level of IgE in patients with severe psoriasis who were not responding to the common therapy.

RESULTS: Patients with severe psoriasis had high levels of IgE.

CONCLUSIONS: IgEs have an important role in pathogenesis of psoriasis and atopic dermatitis, but the mechanism is not yet clear.








IHC_CD34









Immunohistochemical study of osteopontin, Ki-67, and CD34 of psoriasis in Mansoura, Egypt, Maha Mohamed Amin *et al*, 2012



High Magnification




x400



IHC_Loricrin (Skin barrier)





IHC_S100A9 (Inflammation maker)



S100A9
Normal

S100A9
IMQ



S100A9
Plasma

S100A9
IMQ+Plasma

