

Accepted Manuscript

Low-intensity current (LIC) stimulation of subcutaneous adipose derived stem cells (ADSCs) - A missing link in the course of LIC based wound healing


Ravindra V. Badhe, Sonali S. Nipate

PII: S0306-9877(18)31280-5
DOI: <https://doi.org/10.1016/j.mehy.2019.02.039>
Reference: YMEHY 9148

To appear in: *Medical Hypotheses*

Received Date: 14 December 2018
Revised Date: 13 February 2019
Accepted Date: 15 February 2019

Please cite this article as: R.V. Badhe, S.S. Nipate, Low-intensity current (LIC) stimulation of subcutaneous adipose derived stem cells (ADSCs) - A missing link in the course of LIC based wound healing, *Medical Hypotheses* (2019), doi: <https://doi.org/10.1016/j.mehy.2019.02.039>



This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Low-intensity current (LIC) stimulation of subcutaneous adipose derived stem cells (ADSCs) - A missing link in the course of LIC based wound healing

Ravindra V. Badhe, PhD^{1*}

Sonali S. Nipate, PhD²

¹. Dr. D. Y. Patil Institute of Pharmaceutical Sciences and Research, Pimpri, Pune, Maharashtra, India.

². PE Society's Modern College of Pharmacy, Nigdi, Pune, Maharashtra, India.

*Contact Information:

Ravindra V. Badhe

Phone No. +91 9422432038

Email ID. ravindra.badhe@dypvp.edu.in

ABSTRACT

Millions of people die as a result of fatal injuries accounting for 9% of the total global annual deaths. Non fatal injuries generally result in variety of wounds. The normal wound healing process is slow and takes weeks to months, depending on the type of wound. In last two decades, electrotherapy called Low-intensity currents (LIC) for the treatment became popular for faster wound healing, as well as in management of nonresponding and ulcerative wounds. It was reported that LIC mimics 'the current of injury' which is generated by body on wounding and helps in faster wound healing. Researchers have also studied the migration of localized cell and other bio-molecules under the influence of LIC helping the wound to heal faster. Literature review has also suggested that, electrical stimulation of isolated adipose tissue derived stem cells (ADSCs) releases growth factors and differentiates in to specialized cells like fibroblasts and keratinocytes in laboratory conditions. These research areas are well explored and emerged as independent state-of-the-arts therapies and technologies.

Considering the fact, that adipose tissue (along with ADSCs) is present subcutaneously, a new hypothesis is proposed which states that 'Low intensity current (LIC) stimulation of wound stimulates subcutaneous adipose tissue containing ADSCs which releases different growth factors and also differentiates into certain cells like fibroblasts, neurons and keratinocytes. These cells easily migrate to wound site due to lipolysis and loosening of fat tissue, resulting in faster wound healing'. Thus this hypothesis provides a missing link between two state of the art technologies; first one is 'LIC based electrotherapy' and second one is '*in-vitro* LIC stimulation of ADCSs' where role and significance of *in-situ* ADCSs were never studied.

Keywords: Small electric current, adipose tissue derived stem cells (ADSCs), growth factors, wound healing, conducting polymers.

Abbreviation:

LIC - Low intensity current, ADSCs - Adipose Derived Stem Cells, GF - Growth Factors, VEGF - Vascular Endothelial Growth Factors,

INTRODUCTION

An injury that causes a wound is either in the form of a cut, a blow or other impact to living tissue, typically to the skin. According to WHO, Injuries are a global public health problem. More than 5 million people die each year globally accounting for 9% of the world's deaths as a result of injuries. These deaths represent only a small fraction of those injured. Whereas, billions of people suffer from injuries that lead to hospitalization, accident and emergency department, general practitioner treatment or treatment at home ^[1]. Typically there are five types of wounds abrasion, laceration, puncture, avulsion and burn. There are typically six stages of wound healing haemostasis, inflammation, proliferation, angiogenesis, re-epithelialisation and maturation / remodelling ^[2]. The normal healing process is slow and takes weeks to months, depending on type of wound. Nonresponding wounds and ulcers are the most challenging wounds in the wound healing therapy. In this article, new hypothesis for LIC based wound healing is proposed which is based on the understanding that LIC stimulates ADSCs present naturally in subcutaneous adipose tissue, releasing different GFs and differentiating into different cell types which migrate to wound site and help in faster wound healing.

DISCUSSION

Existing State-of-the-art technologies

1. Electrotherapy based Wound healing - In last two decades, many researchers have studied the healing effect of Low-intensity currents (LIC) in the treatment of acute, chronic or nonresponding standard wounds, arterial ulcers, diabetic ulcers, stasis ulcers etc. The results obtained from these studies were very encouraging and thus LIC has gained popularity in the last few years ^[3-11]. LIC mimics 'the current of injury' which is generated by body on wounding. Researchers have proposed that 'the current of injury' is mediated through cell membrane potential and generated by transportation of Na^+ into the cell creating negative pole outside the cell, this system is called as 'Skin battery' ^[12]. The current of injury ranges from 200–800 μA and has been observed to be decreasing in nonresponding wounds. This observation triggered the thought of applying the LIC equivalent to 'current of injury' externally, to improve wound healing (Fig. 1) ^[13].

The mechanism proposed for faster wound healing in this state-of-the-art states that LIC stimulates the migration of the localized cells (WBC, endothelial cells and fibroblasts) in the wound due to membrane potential difference generated by electrodes of LIC device (from + to - terminal). LIC

application to these cells also stimulates release of GFs and collagen at the wound site helping the wound to heal faster.




Figure 1. Known electrotherapy based wound healing mechanism

2. *In-vitro* LIC stimulation of ADSCs - In human body, adipose tissue is mainly located beneath the skin [subcutaneous fat / white adipose tissue (WAT)]. It is also present around internal organs (visceral fat), intermuscular (Muscular system), in bone marrow (yellow bone marrow) and in the breast tissue. It was reported that adipose tissue derived stem cells (ADSCs) obtained from adult adipose tissue provide pool of multi-potent adult stem cells that are noncontroversial in nature, relatively abundant, easy to isolate and are expandable ^[14]. There are recent reports stating electrical stimulations of adipose-derived stem cells (ADSCs) generating specialized differentiated cells like nerve cells, cardiomyocytes, fibroblast, etc. and also stimulating secretion of FGs like VEGF-A, NGF etc. (Fig. 2) ^[15-17].

Thus, this state-of-the-art work proved that LIC stimulation of ADSCs can differentiate them into various functionalised cells and also improves release of GFs. Interestingly; all the studies reported for this type of work are *in-vitro* studies with isolated ADSCs.




Figure 2. Differentiation of Adipose Derived Stem Cells (ADSCs) in the presence of LIC

3. Neurogenesis and Angiogenesis go hand in hand - Neurogenesis and Angiogenesis are critical aspects of wound healing. Insufficient angiogenesis results in impaired wound healing and formation of chronic wound. Recent reports suggest that neurogenesis and angiogenesis share some molecular signals that act synergistically^[18-20]. The expression of vascular endothelial growth factor (VEGF-A) with LIC stimulation of endothelial cells was reported along with the role of other growth factors like Nerve Growth Factor (NGF) in neurogenesis and angiogenesis^[21-23]. It was reported that wounded neurons in wound show nerve regeneration in the direction of LIC stimulation (+ to -). This regeneration takes place mainly due to release of VEGF. This GF is released due to LIC stimulation of Schwann cells present on the wounded neurons (Fig. 3). This increased release of VEGF additionally helps in sprouting of new blood vessels. Thus, LIC induces neurogenesis and angiogenesis, helping wound to heal faster^[24-26].




Figure 3. LIC stimulation of Schwann cells on wounded neurons help in regeneration of neurons and blood vessels.

The literature survey suggests that, LIC positively plays a role in conditioning of isolated ADSCs, differentiating them into specific cell types and secretion of certain growth factors. Improved wound healing due to LIC was also well established. However, there are no studies reported on the effect of LIC stimulation of ADSCs present *in-situ* in adipose tissue. Thus, there is still a 'gap' between these two State-of-the-art research areas. This gap has provided the unique opportunity to explore the effect of LIC stimulation of subcutaneous adipose tissue containing ADSCs specifically from the wound healing prospective.

Hypothesis

By understanding the gap between the present studies and established technologies and taking advantage of the opportunity to explore new research area, a novel hypothesis is proposed here stating 'low intensity current (LIC) stimulation of wound stimulates subcutaneous adipose tissue containing ADSCs which releases different growth factors and also differentiates into certain cells like fibroblasts, neurons and keratinocytes. These cells easily migrate to wound site due to lipolysis and loosening of fat tissue, resulting in faster wound healing'.

Probable wound healing mechanism of proposed hypothesis

The LIC wound healing therapy is based on two key components one is hydrogel and other is electrotherapy device. Hydrogel when applied on the wound fills the wound completely and uniformly. It provides the moist environment for wound healing and distributes LIC uniformly throughout the wound. The proposed hypothesis predict that the applied electric current passes through hydrogel and wound surface stimulating WBC, endothelial cells, fibroblasts and shwann cells present on the damaged neurons. Stimulation of endothelial cells and fibroblast release VEGF and collagen respectively followed by migration and accumulation of collagen and fibroblast cells at wound site. Stimulation of shwann cells on neuron release VEGF helping in sprouting and regenerating neurons and blood vessels. Further, when electric current reaches the adipose tissue, two phenomenon take place. First the LIC will lyses (lipolysis) some adipocyte cells opening the tight junction between the adipocyte cells and loosening fat tissue. This will help in faster migration of growth factors and differentiated cells generated by LIC stimulation of ADSCs to the wound site, which is the second phenomenon. Thus, if we combine all the effects of LIC stimulation of the wound, overall we get faster wound healing which is the likely mechanism for proposed hypothesis (Fig. 4) ^[27]. The novelty of the hypothesis lies in bridging the gap between electrotherapy based wound healing studies and isolated ADSCs based wound healing studies.




Figure 4. Mechanism of Proposed Hypothesis

Upshot of the hypothesis - Advancement and innovations in science.

1. Understanding the role of swelled or inflamed subcutaneous fat cells on regulation of ADSCs and wound healing'

The hypothesis will support the claim of faster wound healing owing to the LIC stimulation of subcutaneous ADSCs leading to differentiation of the stem cells and growth factor release. All the normal healthy individuals have some amount of adipose tissue preset subcutaneously, thus we expect improved wound healing with LIC stimulation in these individuals.

It is worth to study the effect of LIC stimulation on wound healing in obese individuals. This study will be very interesting as there are two contradictory conditions. Obese individual will have excess amount of adipose tissue developed due to adipose homeostasis associated with blocked insulin signals with inflammatory cytokines released by inflamed adipose tissue ^[28]. Thus, these individuals will have comparatively more number of ADSCs and will help in faster wound healing. On the other hand, the swelled or inflamed adipose tissue will tightly enclose ADSCs in the excess of adipocytes cells which are full of fat. The presence of excess of tightly arranged [very less water filled extracellular matrix (ECM)] swelled adipose cells (full of fat which is poor conductor of electricity) might increase the resistance to the LIC and further hamper wound healing ^[29].

Hence it will be interesting to understand the effect of LIC on modulation of up-regulation and down-regulation of ADSCs in normal and obese individuals. It will also shade a light on effect of

LIC on modulation of adipose homeostasis and insulin resistance. Thus both of these areas can become innovations in science.

2. Developing advanced wound healing therapies

As mentioned earlier, LIC based wound healing therapy is based on two components, hydrogel and electrotherapy device. Hydrogel when applied on the wound fills the wound completely and uniformly, helping to distribute the LIC equally through the wound. The formulation of hydrogel itself is an established technology and developing hydrogel for LIC based wound healing will take it to the next level. The advancement is possible by formulating medicated hydrogel. Here, hydrogel containing pain relieving medicament, antimicrobial agents, nutrients and growth factors/promoters can be formulated to improve wound healing ^[30-36]. One can improve the electrical conducting property of the hydrogel using metallic nanoparticles and conducting polymers ^[37-40] or a medicated conducting hydrogel combining both the properties can also be formulated and used to improve LIC based wound healing. The electrotherapy device can also be advanced by developing handheld unit with varying current (from 10 - 800 μ A) and flexible moving electrodes. These advancements in present LIC based wound healing therapies in the light of proposed hypothesis might enhance wound healing therapy significantly.

CONCLUSION

In conclusion, the thorough understanding of present state of LIC based wound healing therapies, LIC stimulation experiments on isolated ADSCs and LIC based neurogenesis and angiogenesis experiments suggested faster wound healing with LIC but these studies evolved separately. The proposed hypothesis in this article suggest the cohere effect of all the above mentioned studies systematically playing individual roles for rapid wound healing.

The hypothesis states that 'low intensity current (LIC) stimulation of wound stimulates subcutaneous adipose tissue containing ADSCs which releases different growth factors, and also differentiates into certain cells like fibroblasts, neurons and keratinocytes, which migrate easily to wound site due to lipolysis and loosening of fat tissue, resulting in faster wound healing'. The probable mechanism showing the cohere action of individual studies reported as 'the applied electric current passes through wound stimulating WBC, endothelial cells, fibroblasts and shwann cells present on the damaged neuron. Stimulation of endothelial cells and fibroblast release VEGFs and collagen respectively followed by deposition of the same on wound site, stimulation of shwann cells on neuron also release VEGF helping in sprouting and regenerating neurons and blood vessels. As

the electric current reaches the adipose tissue, two phenomenon occur. First the LIC will lyses (lipolysis) some adipocyte cells opening the tight junction between the adipocyte cells and leads to loosening of fat tissue. This will help in faster migration of growth factors and differentiated cells generated by LIC stimulation of ADSCs (which is the second phenomenon which occur simultaneously) at the wound site helping in faster wound healing.

The proposed hypothesis can advance the present LIC based wound healing therapy to next level using medicated, conductive (improved) hydrogel and handheld electrotherapy device. It will also provide the knowledge about the role of swelled or inflamed subcutaneous fat cells on regulation of ADSCs and ultimately in wound healing of obese / diabetic obese people.

CONFLICT OF INTEREST

Non

FUNDING

Funded Internally

REFERENCES

- [1] WHO report - Injuries and Violence the Facts - 2014
(<http://www.who.int/iris/handle/10665/149798>)
- [2] Velnar T, Bailey T, and Smrkolj V. The Wound Healing Process: an Overview of the Cellular and Molecular Mechanisms. *The Journal of International Medical Research*. 2009; 37: 1528 – 1542.
- [3] Yadollahpour A, Rashidi S, and Fayazi R. Electrical Stimulation for Chronic Wound Treatment: Basic Principles and Current Clinical Position. *J Med Res Prac*. 2017; 6(3): 76–80.
- [4] Ud-Din S and Bayat A. Electrical Stimulation and Cutaneous Wound Healing: A Review of Clinical Evidence. *Healthcare (Basel)*. 2014; 2(4): 445–467.

- [5] Ashrafi M, Alonso-Rasgado T, Baguneid M, Bayat A. The efficacy of electrical stimulation in experimentally induced cutaneous wounds in animals. *Vet Dermatol*. 2016; 27(4): 235-e57.
- [6] Tai G, Tai M and Zhao M. Electrically stimulated cell migration and its contribution to wound healing. *Burns Trauma*. 2018; 6: 20.
- [7] Iwasa SN, Babona-Pilipos R, Morshead CM. Environmental Factors That Influence Stem Cell Migration: An "Electric Field". *Stem Cells Int*. 2017, 2017, 4276927.
- [8] Simpson MJ, Lo KY, Sun YS. Quantifying the roles of random motility and directed motility using advection-diffusion theory for a 3T3 fibroblast cell migration assay stimulated with an electric field. *BMC Syst Biol*. 2017, 11(1), 39.
- [9] Polak A, Kucio C, Kloth LC. A randomized, controlled clinical study to assess the effect of anodal and cathodal electrical stimulation on periwound skin blood flow and pressure ulcer size reduction in persons with neurological injuries. *Ostomy Wound Manage*. 2018; 64(2):10-29.
- [10] Khouri C, Kotzki S, Roustit M, Blaise S, Gueyffier F, Cracowski JL. Hierarchical evaluation of electrical stimulation protocols for chronic wound healing: An effect size meta-analysis. *Wound Repair Regen*. 2017; 25(5): 883-891.
- [11] Houghton PE. Electrical stimulation therapy to promote healing of chronic wounds: a review of reviews. *Chronic Wound Care Research and Management*. 2017; 2017(4): 25-44.
- [12] Barker AT, Jaffe LF, and Vanable JW. The glabrous epidermis of cavies contains a powerful battery. *Am J Physiol Regulatory Integrative Comp Physiol*. 1982; 242: 358-366.
- [13] Balakatounis KC and Angoules AG. Low-intensity Electrical Stimulation in Wound Healing: Review of the Efficacy of Externally Applied Currents Resembling the Current of Injury. *Open Access Journal of Plastic Surgery*. 2008; 8: 283-291.

- [14] Pavese A, Soncini M, Zamperone A, Pietronave S, Medico E, Redaelli A, Prat M, Fiore GB, Electrical Conditioning of Adipose-Derived Stem Cells in a Multi-Chamber Culture Platform. *Biotechnology and Bioengineering*. 2014; 111(7): 1452-1463
- [15] Wankhade UD, Shen M, Kolhe R, Fulzele S. Advances in Adipose-Derived Stem Cells Isolation, Characterization, and Application in Regenerative Tissue Engineering. *Stem Cells International*. 2016: Article ID 3206807.
- [16] Tandon N, Goh B, Marsano A, Chao PHG, Sorrentino CM, Gimble J, and Novakovic GV. Alignment and Elongation of Human Adipose-Derived Stem Cells in Response to Direct-Current Electrical Stimulation. *Conf Proc IEEE Eng Med Biol Soc*. 2009; 1: 6517–6521.
- [17] Yang Y, Ma T, Ge J, Quan X, Yang L, Zhu S, Huang L, Liu Z, Liu L, Geng D, Huang J, and Luo Z. Facilitated Neural Differentiation of Adipose Tissue–Derived Stem Cells by Electrical Stimulation and Nurr-1 Gene Transduction. *Cell Transplantation*. 2016; 25: 1177–1191.
- [18] Huang J, Zhang Y, Lu L, Hu X and Luo Z. Electrical stimulation accelerates nerve regeneration and functional recovery in delayed peripheral nerve injury in rats. *European Journal of Neuroscience*. 2013; 38: 3691–3701.
- [19] Ud-Din S, Sebastian A, Giddings P, Colthurst J, Whiteside S, Morris J, Nuccitelli R, Pullar C, Baguneid M, Bayat A. Angiogenesis Is Induced and Wound Size Is Reduced by Electrical Stimulation in an Acute Wound Healing Model in Human Skin. *PLOS ONE*. 2015; 10(4): e0124502.
- [20] Lin YC, Kao CH, Chen CC, Ke CJ, Yao CH, Chen YS. Time-Course Effect of Electrical Stimulation on Nerve Regeneration of Diabetic Rats. *PLOS ONE*. 2015; 10(2): e0116711
- [21] Liebano RE. and Machado AFP., Vascular Endothelial Growth Factor Release Following Electrical Stimulation in Human Subjects. *Advances In Wound Care*. 2013; 3(2): 98-103.

- [22] Wild R, Klems A, Takamiya M, Hayashi Y, Strahle U, Ando K, Mochizuki N, van Impel A, Schulte-Merker S, Krueger J, Preau L, le Noble F. Neuronal sFlt1 and VEGFA determine venous sprouting and spinal cord vascularization. *Nature Communications*. 2017; 8: 13991.
- [23] Al-Majed AA, Brushart TM, Gordon T. Electrical stimulation accelerates and increases expression of BDNF and trkB mRNA in regenerating rat femoral motoneurons. *Eur J Neurosci*. 2000; 12: 4381–4390.
- [24] Mackenzie F and Ruhrberg C, Diverse roles for VEGF-A in the nervous system. *Development*. 2012; 139: 1371-1380.
- [25] McCaig CD, Song B and Rajnicek AM, Electrical dimensions in cell science. *Journal of Cell Science*. 2009; 122(23): 4267-4276.
- [26] Himmels P, Paredes I, Adler H, Karakatsani A, Luck R, Marti HH, Ermakova O, Rempel E, Stoeckli ET, de Almodóvar CR. Motor neurons control blood vessel patterning in the developing spinal cord. *Nature Communications*. 2017; 8: 14583.
- [27] Hamida ZH, Comtois AS, Portmann M, Boucher JP, Savard R, Effect of electrical stimulation on lipolysis of human white adipocytes. *Appl. Physiol. Nutr. Metab*. 2011; 36: 271–275.
- [28] Borsta SE, Conover CF. High-fat diet induces increased tissue expression of TNF- α . *Life Sciences*. 2005; 77: 2156–2165.
- [29] Miklavcic D, Pavselj N, Hart FX. Electric Properties of Tissues. *Wiley Encyclopedia of Biomedical Engineering*. 2006 (Conference Proceedings).
- [30] Roy N, Saha N, Kitano T, Saha P, Development and Characterization of Novel Medicated Hydrogels for Wound Dressing. *Soft Materials*. 2010; 8(2): 130-148.

- [31] Badhe RV, Nanda RK, Chejara DR, Choonara YE, Kumar P, du Toit LC, Pillay V, Microwave-assisted facile synthesis of a new tri-block chitosanconjugate with improved mucoadhesion. *Carbohydrate Polymers*. 2015; 130: 213–221.
- [32] Badhe RV., Bijukumar D., Chejara DR., Mabrouk M, Choonara YE., Kumar P., du Toit LC., Kondiah PPD., Pillay V. A composite chitosan-gelatin bi-layered, biomimetic macroporous scaffold for blood vessel tissue engineering. *Carbohydrate Polymers*. 2017; 157: 1215–1225.
- [33] Griffin DR, Weaver WM, Scumpia PO, Di Carlo D, Segura T. Accelerated wound healing by injectable microporous gel scaffolds assembled from annealed building blocks. *Nat Mater*. 2015; 14(7): 737–744.
- [34] Liu H, Wang C, Li C, Qin Y, Wang Z, Yang F, Li Z, and Wang J. A functional chitosan-based hydrogel as a wound dressing and drug delivery system in the treatment of wound healing. *RSC Adv.*, 2018; 8: 7533-7549.
- [35] Zhu J, Han H, Ye T-T, Li F-X, Wang X-L, Yu J-Y, and Wu D-Q. Biodegradable and pH Sensitive Peptide Based Hydrogel as Controlled Release System for Antibacterial Wound Dressing Application. *Molecules*. 2018; 23: 3383.
- [36] Mir M, Ali MN, Barakullah A, Gulzar A, Arshad M, Fatima S and Asad M. Synthetic polymeric biomaterials for wound healing: a review. *Prog Biomater*. 2018; 7(1):1-21.
- [37] Mawad D, Lauto A., Wallace GG. Conductive Polymer Hydrogels in Polymeric Hydrogels as Smart Biomaterials, S. Kalia (ed.), Polymeric Hydrogels as Smart Biomaterials, Springer Series on Polymer and Composite Materials. Springer International Publishing, Switzerland, 2016, pp. 19-44.
- [38] Thoniyot P, Tan MJ, Karim AA, James D, Xian Y, Loh J. Nanoparticle–Hydrogel Composites: Concept, Design, and Applications of These Promising, Multi- Functional Materials. *Adv.Sci*. 2015; 2: 1400010.

- [39] Alonso E, Faria M, Mohammadkazemi F, Resnik M, Ferreira A, Cordeiro N. Conductive bacterial cellulose-polyaniline blends: Influence of the matrix and synthesis conditions. *Carbohydrate Polymers*. 2018; 183:254-262.
- [40] Zarrintaj P, Rezaeian I, Bakhshandeh B, Heshmatian B, Ganjali MR. Bio - Conductive Scaffold Based on Agarose - Polyaniline for Tissue Engineering, *J Skin Stem Cell*. 2017; 4(2): E67394.

ACCEPTED MANUSCRIPT